

2XYT Growth Medium

Intended Use

2XYT Growth Medium is an optimized formulation for the growth and maintenance of M13 phage or other filamentous ss DNA bacteriophages.

Summary

2XYT Growth Medium was originally formulated as a nutritionally enriched growth medium for growth of recombinant strains of *Escherichia coli* and can also be used for propagation of M13 bacteriophage. It permits larger quantity of phage production without exhausting the host. This media is 2 times richer than the YT media.

Principle

Yeast extract and Tryptone provide all the required amino acids, nucleotide precursors, vitamins and other metabolites and as a result the cells grow faster in this medium. Sodium chloride provides sodium ions for transport and osmotic balance.

Formula*

Ingredients	g/L
Tryptone	16.0
Yeast Extract	10.0
Sodium Chloride	5.0
Final pH (at 25°C)	7.0 ± 0.2

*Adjusted to suit performance parameters.

Storage and Stability

Store dehydrated medium below 30°C in tightly closed container and the prepared medium at 2°C-8°C. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Specimen Collection and Handling

Ensure that all samples are properly labelled. Follow appropriate techniques for handling samples as per established guidelines. Some samples may require special handling, such as immediate refrigeration or protection from light, follow the standard procedure. The samples must be stored and tested within the permissible time duration. After use, contaminated materials must be sterilized by autoclaving before discarding.

Directions

1. Suspend 31.00 g of the powder in 1000 mL purified / distilled water.
2. Heat to boiling to dissolve the powder completely.
3. Dispense as desired and sterilize by autoclaving at 121°C (15 psi) for 15 minutes as per validated cycle.

Quality Control

Dehydrated Appearance: Yellow to beige, homogeneous free flowing powder.

Prepared Appearance: Light yellow to yellow coloured, clear solution without any precipitate.

Growth Promotion Test: Growth promotion is carried out in accordance with the harmonized method of USP/EP/JP/IP and growth is observed after an incubation at 30°C-35°C for 18 to 24 hours.

Growth Promoting Properties: The test results observed are within the specified temperature and shortest period of time specified in the test, inoculating ≤ 100 cfu of appropriate microorganism at 30°C-35°C for 18 hours.

Organism (ATCC)	Growth
<i>Escherichia coli</i> (23724)	Good
<i>Escherichia coli</i> (53868)	Good

Performance and Evaluation

Performance of the product is dependent on following parameters as per product label claim:

1. Directions
2. Storage
3. Expiry

Warranty

This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

Reference

1. Difco manual 11th ed., Sparks, MD (1998), 22-23
2. Assubel, F.M., R. Brent, R.E. Kingston, D.D. Moore, J.G. Seidman, J.A. Smith and K. Struhl, Current protocols in molecular biology, vol. 1, Current Protocols, New York, (1994)
3. Davis, L.G., M.D. Dibner and J.F. Battey, Basic methods in molecular biology, Elsevier, New York, (1986).
4. Data on file: Microxpress®, A Division of Tulip Diagnostics (P) Ltd.

Product Presentation:

Cat. No.	Product Description	Pack Size
201270050500	Dehydrated Culture Media	500 g

 Temperature Limit	 Manufacturer	 LOT	Batch Code	 Date of Manufacture	 This way up	 RO Received on
REF Catalogue Number	 Consult Instructions for use	 Use-by Date	 HC	Hygroscopic keep container tightly closed	OO Opened on	

Revision: 0825/VER-03

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.