

## Buffered Peptone Water

### Intended Use

Buffered Peptone Water used for pre-enrichment of injured *Salmonella* species. It increases recovery of *Salmonella* species prior to selective enrichment from food.

### Summary

Buffered Peptone Water (BPW) is one of the most widely used pre-enrichment broths for *Salmonella* in a wide range of foods. Moreover, it is often the medium of choice in many published reference methods, including the current International Organization for Standardization (ISO) method, for detection of *Salmonella* in foods.

Edel and Kampelmaacher in their comparative study observed that food preservation methods cause sublethal injury to *Salmonellae*. It is known that freezing and drying may injure *Salmonellae* so that they are unable to multiply in selective media.

Pre-enrichment of sublethally injured *Salmonellae* in buffered peptone water showed superior results in comparison with direct selection methods. In addition to providing conditions for the recovery and growth of cells prior to selective enrichment, BPW buffers the pH of the growth system against pH changes brought about by the growth and metabolism of microorganisms during enrichment and those imposed by the food sample.

### Principle

Proteose peptone serves as a source of carbon, nitrogen, vitamins and minerals. Sodium chloride provides sodium ions for the membrane transport and maintains osmotic equilibrium of the medium. Phosphates buffer the medium.

### Formula\*

Ingredients	g/L
Proteose Peptone	10.0
Sodium Chloride	5.0
Disodium Phosphate	3.5**
Monopotassium Phosphate	1.5
Final pH (at 25°C)	7.2 ± 0.2

\*Adjusted to suit performance parameters.

\*\*3.5 g of Disodium Phosphate, Anhydrous is equivalent to 9.0 g of Disodium Phosphate Dodecahydrate.

### Storage and Stability

Store dehydrated medium below 30°C in tightly closed container and the prepared medium at 2°C-8°C. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

### Type of specimen

Food and dairy samples.

### Specimen Collection and Handling

Ensure that all samples are properly labelled.

Follow appropriate techniques for handling samples as per established guidelines.

Some samples may require special handling, such as immediate refrigeration or protection from light, follow the standard procedure.

The samples must be stored and tested within the permissible time duration.

After use, contaminated materials must be sterilized by autoclaving before discarding.

### Directions

1. Suspend the 20.00 g (Equivalent weight of dehydrated medium) of the powder in 1000 mL purified / distilled water.
2. Mix thoroughly and warm slightly with frequent agitation to dissolve the powder completely.
3. Sterilize by autoclaving at 121°C (15 psi) for 15 minutes as per validated cycle.

## Quality Control

**Dehydrated Appearance:** Cream to yellow coloured, homogenous, free flowing powder.

**Prepared Appearance:** Straw yellow to light yellow coloured, clear solution without any precipitate.

**Growth Promotion Test:** Growth promotion is carried out in accordance with the harmonized method of USP/EP/JP/IP and growth is observed after an incubation at 30°C-35°C for 18-24 hours.

**Growth Promoting Properties:** The test results observed are within the specified temperature and shortest period of time specified in the test, inoculating ≤ 100 cfu of appropriate microorganism at 30°C-35°C for 18 hours.

Organisms (ATCC)	Growth
<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Abony</i> (NCTC 6017)	Good
<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Typhimurium</i> (14028)	Good
<i>Escherichia coli</i> (8739)	Good

## Interpretation of Results

Growth in the medium is indicated by the presence of turbidity compared to an uninoculated control.

## Performance and Evaluation

Performance of the product is dependent on following parameters as per product label claim:

1. Directions
2. Storage
3. Expiry

## Warranty

This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

## Reference

1. Edel and Kampelmacher, 1973, Bull. W.H.O., 48:167.
2. Sadovski, 1977, J. Food Technol., 12:85.
3. Juven, Cox, Bailey, Thomson, Charles and Schutze, 1984, J. Food Prot., 47:299
4. Data on file: Microxpress®, A Division of Tulip Diagnostics (P) Ltd.

## Product Presentation:

Cat No.	Product description	Pack Size
201020370100	Dehydrated Culture Media	100 g
201020370500	Dehydrated Culture Media	500 g

 Temperature Limit	 Manufacturer	 <b>LOT</b>	 Batch Code	 Date of Manufacture	 This way up	 Received on
<b>REF</b> Catalogue Number	 Consult Instructions for use	 Use-by Date	 Hygroscopic keep container tightly closed	<b>OO</b> Opened on		

Revision: 0725/VER-03

## Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.