

Lactose Monohydrate Broth (Broth Medium D) BP

Intended Use

Lactose Monohydrate Broth (Broth Medium D) is used as a medium for the detection of coliforms, in water, food and dairy products in accordance with British Pharmacopoeia.

Summary

Lactose Broth is recommended by British pharmacopoeia for selective pre-enrichment of *Enterobacteriaceae* as well as for *E. coli* and *Salmonella* in water, food and pharmaceutical products. The medium is used for detection of specified microorganisms of non-sterile products according to British Pharmacopoeia. This medium is also recommended by various other pharmacopoeia. Lactose Monohydrate Broth is recommended by APHA in the performance and confirmation of the presumptive test for coliform bacteria in water, food and milk. This medium can be used as an alternate to Buffered Sodium Chloride-peptone Solution pH 7.0 for the revival of *Enterobacteriaceae* and Gram-negative bacteria. The medium is incubated for a time sufficient to revive the bacteria but not the multiplication of the bacteria. It is recommended to incubate the medium usually for 2 hours and not for more than 5 hours.

Principle

Pancreatic digest of gelatin and cara beef extract supply essential nutrients to the organisms. Lactose monohydrate is a fermentable carbohydrate.

Formula*

Ingredients	g/L
Pancreatic Digest of Gelatin	5.0
Cara Beef Extract [#]	3.0
Lactose Monohydrate	5.0
Final pH (at 25°C)	6.9 ± 0.2

*Adjusted to suit performance parameters.

[#]Equivalent to Beef Extract

Storage and Stability

Store dehydrated medium below 30°C in tightly closed container and the prepared medium at 2°C-8°C. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Type of specimen

Food and Dairy samples; Water samples

Specimen Collection and Handling

Ensure that all samples are properly labelled.

Follow appropriate techniques for handling samples as per established guidelines.

Some samples may require special handling, such as immediate refrigeration or protection from light, follow the standard procedure.

The samples must be stored and tested within the permissible time duration.

After use, contaminated materials must be sterilized by autoclaving before discarding.

Directions

1. Suspend 12.75 g of the powder in 1000 mL purified/distilled water.
2. Heat if necessary, to dissolve the powder completely.
3. For larger inocula (10 mL or more), concentrated medium may be prepared to account for medium dilution by the inoculum.
4. Dispense in tubes containing inverted fermentation vial (Durham's tube) as desired.
5. Sterilize by autoclaving at 121°C (15 psi) for 15 minutes as per validated cycle and cool immediately.

Quality Control

Dehydrated Appearance: Yellow coloured, homogeneous, free flowing powder.

Prepared Appearance: Light yellow to medium amber coloured, clear solution without any precipitate.

Cultural Response: Growth is observed after an incubation at 30°C -35°C for 18-48 hours.

Organisms (ATCC)	Growth	Gas	Acid
<i>Klebsiella aerogenes</i> (13048)	Good	+	+
<i>Enterococcus faecalis</i> (29212)	Good	-	-
<i>Escherichia coli</i> (8739)	Good	+	+
<i>Escherichia coli</i> (25922)	Good	+	+
<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Typhimurium</i> (14028)	Good	-	-

Note:

After incubation, add 1-2 drops of 1% phenol red solution to observe acid production.

Performance and Evaluation

Performance of the product is dependent on following parameters as per product label claim:

1. Directions
2. Storage
3. Expiry

Warranty

This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

Reference

1. Eaton A. D., Clesceri L. S. and Greenberg A W (Eds.), 2008, Standard Methods for the Examination of Water and Wastewater, 21st ed., APHA, Washington, D.C.
2. Downes F P and Ito K(Eds.), 2001, Compendium of Methods for The Microbiological Examination of Foods, 4th ed., APHA, Washington, D.C
3. Wehr H M and Frank J H., 2004, Standard Methods for the Examination of Dairy Products, 17th ed., APHA Inc., Washington, D.C.
4. British Pharmacopoeia, 2009, The Stationery office British Pharmacopoeia.
5. Data on file: Microxpress®, A Division of Tulip Diagnostics (P) Ltd.

Product Presentation:

Cat No.	Product description	Pack Size
201120070100	Dehydrated Culture Media	100 g
201120070500	Dehydrated Culture Media	500 g

 Temperature Limit	 Manufacturer	 LOT	 Batch Code	 Date of Manufacture	 This way up	 Received on
REF Catalogue Number	 Consult Instructions for use	 Use-by Date	 Hygroscopic keep container tightly closed	OO Opened on		

Revision: 0825/VER-03

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.