

Moeller Decarboxylase Broth with Ornithine

Intended Use

Moeller Decarboxylase Broth with Ornithine is used to differentiate bacteria on the basis of their ability to decarboxylate L-Ornithine.

Summary

Moeller Decarboxylase Broth with Ornithine hydrochloride is used for differentiating Gram-negative enteric bacilli on the basis of their ability to decarboxylate L-Ornithine. Decarboxylase Broth was introduced by Moeller for detecting the production of lysine and ornithine decarboxylase and arginine dihydrolase. Prior to Moeller's work, bacterial amino acid decarboxylases were studied by Gale, and Gale and Epps. Production of ornithine decarboxylase is helpful criterion in differentiating *Klebsiella* and *Enterobacter* species. *Klebsiella* are nonmotile and do not produce ornithine decarboxylase while *Enterobacter* are motile and produce ornithine decarboxylase except *Enterobacter agglomerans*. Decarboxylase media are also recommended by standard methods for identification of bacteria.

Principle

This medium contains beef extract and peptic digest of animal tissue which provide nitrogenous nutrients for the growth of bacteria. Dextrose is the fermentable carbohydrate and pyridoxal is the co-factor for the decarboxylase enzyme. Bromocresol purple and cresol red are the pH indicators in this medium. When the medium is inoculated with the dextrose fermenting bacteria, the pH is lowered due to acid production which changes the colour of the indicator from purple to yellow. Acid produced stimulates decarboxylase enzyme. Putrescine is produced due to ornithine decarboxylation. Formation of amine putrescine increases the pH of the medium, changing the colour of the indicator from yellow to purple. If the organisms do not produce the appropriate enzyme, the medium remains acidic, yellow in colour.

Formula*

Ingredients	g/L
Peptic Digest of Animal Tissue	5.0
Beef Extract	5.0
Dextrose	0.5
Bromocresol Purple	0.01
Cresol Red	0.005
Pyridoxal	0.005
L-Ornithine	10.0
Final pH (at 25°C)	6.0 ± 0.2

*Adjusted to suit performance parameters.

Storage and Stability

Store dehydrated medium below 30°C in tightly closed container and the prepared medium at 2°C-8°C. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Specimen Collection and Handling

Ensure that all samples are properly labelled. Follow appropriate techniques for handling samples as per established guidelines. Some samples may require special handling, such as immediate refrigeration or protection from light, follow the standard procedure. The samples must be stored and tested within the permissible time duration. After use, contaminated materials must be sterilized by autoclaving before discarding.

Directions

1. Suspend 20.52 g of the powder in 1000 mL purified / distilled water.
2. Heat if necessary, to dissolve the powder completely.
3. Dispense in 5 mL amount in screw-capped tubes and sterilize by autoclaving at 121°C (15 psi) for 10 minutes as per validated cycle.
4. Cool the tubed medium in an upright position.
5. Inoculate the tubes and overlay with 2-3 mL of sterile mineral oil.

Quality Control

Dehydrated Appearance: Light grey to light green homogeneous free flowing powder.

Prepared Appearance: Purple coloured, clear solution without any precipitate.

Cultural Response: Cultural characteristics observed after an incubation at 35°C-37°C for upto 4 days (Inoculated tubes are overlaid with sterile mineral oil).

Organism (ATCC)	Growth	Ornithine Decarboxylation
<i>Escherichia coli</i> (25922)	Good	±
<i>Klebsiella aerogenes</i> (13048)	Good	+
<i>Klebsiella pneumoniae</i> (13883)	Good	-
<i>Proteus mirabilis</i> (25933)	Good	-
<i>Shigella flexneri</i> serotype 2b (12022)	Good	-
<i>Salmonella Typhi</i> (6539)	Good	-
<i>Citrobacter freundii</i> (8090)	Good	±
<i>Proteus hauseri</i> (13315)	Good	-
<i>Pseudomonas aeruginosa</i> (9027)	Good	-
<i>Salmonella Paratyphi A</i> (9150)	Good	-
<i>Serratia marcescens</i> (8100)	Good	+
<i>Shigella dysenteriae</i> (13313)	Good	-
<i>Shigella sonnei</i> (25931)	Good	+

Key: + = Positive reaction, purple colour

- = Negative reaction, yellow colour

± = Variable (purple / yellow colour)

Interpretation of Results

Each isolate to be tested should also be inoculated into the basal medium tube lacking the amino acid. After incubation, a decarboxylase test may show two layers of different colours, yellow and purple. Shake the tube gently before interpreting the results.

Performance and Evaluation

Performance of the product is dependent on following parameters as per product label claim:

1. Directions
2. Storage
3. Expiry

Precautions / Limitations

Inoculated tubes must be protected from air with a layer of sterile mineral oil. Exposure to air may cause alkalization at the surface of the medium which makes the test invalid.

Warranty

This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

Reference

1. Moeller V., 1955, Acta Pathol. Microbiol. Scand. 36:158.
2. Gale G. F., 1940, Biochem. J., 34:392.
3. Gale and Epps, 1943, Nature, 152:327
4. Data on file: Microxpress®, A Division of Tulip Diagnostics (P) Ltd.

Product Presentation:

Cat No.	Product description	Pack Size
201130630100	Dehydrated Culture Media	100 g

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.
