

Phenol Red Sorbitol Broth

Intended Use

Phenol Red Sorbitol Broth is used for sorbitol fermentation studies of microorganisms.

Summary

Phenol Red Broth Medium is formulated as per Vera and is recommended to determine the fermentation reaction of carbohydrates for the differentiation of microorganisms. Phenol Red Broth Medium with various carbohydrates serves as a differential medium by aiding in differentiation of various species and genera by their ability to ferment the specific carbohydrate, with the production of acid or acid and gas. Phenol Red Sorbitol Broth is used to study Sorbitol fermentation in various bacteria.

Principle

Proteose peptone and cara beef extract serve as sources for carbon and nitrogen. Sodium chloride is the osmotic stabilizer. Phenol red is the pH indicator, which turns yellow at acidic pH i.e. on fermentation of sorbitol. Gas formation is seen in Durham's tubes. All of the *Enterobacteriaceae* grow well in this medium. In addition to producing a pH colour shift, the production of mixed acids, notably butyric acids, often results in a pungent, foul odour from the culture medium.

Formula*

Ingredients	g/L
Proteose Peptone	10.0
Cara Beef Extract [#]	1.0
Sodium Chloride	5.0
Sorbitol	5.0
Phenol Red	0.018
Final pH (at 25°C)	7.4 ± 0.2

*Adjusted to suit performance parameters.

[#] Equivalent to Beef Extract.

Storage and Stability

Store dehydrated medium below 30°C in tightly closed container and the prepared medium at 2°C-8°C. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines.

For food and dairy samples, follow appropriate techniques for handling specimens as per established guidelines.

For water samples, follow appropriate techniques for handling specimens as per established guidelines and local standards.

Specimens should be obtained before antimicrobial agents have been administered.

After use, contaminated materials must be sterilized by autoclaving before discarding.

Directions

1. Suspend 21.01 g of the powder in 1000 mL purified / distilled water.
2. Mix thoroughly.
3. Heat to dissolve the powder completely.
4. Dispense in tubes containing inverted Durham's tubes.
5. Sterilize by autoclaving at 121°C (15 psi) for 15 minutes as per validated cycle.

Quality Control

Dehydrated Appearance: Light yellow to pink coloured, homogeneous, free flowing powder.

Prepared Appearance: Red coloured, clear solution without any precipitate.

Cultural Response: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism (ATCC)	Growth	Acid	Gas
<i>Citrobacter freundii</i> (8090)	Good	Positive reaction, yellow colour	Positive reaction
<i>Escherichia coli</i> (25922)	Good	Positive reaction, yellow colour	Positive reaction
<i>Klebsiella aerogenes</i> (13048)	Good	Positive reaction, yellow colour	Positive reaction
<i>Klebsiella pneumoniae</i> subsp. <i>pneumoniae</i> (10031)	Good	Positive reaction, yellow colour	Positive reaction
<i>Proteus hauseri</i> (13315)	Good	Negative reaction, no colour change	Negative reaction
<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Typhimurium</i> (14028)	Good	Positive reaction, yellow colour	Positive reaction
<i>Shigella flexneri</i> serotype 2b (12022)	Good	Positive reaction, no colour change	Negative reaction

Performance and Evaluation

Performance of the product is dependent on following parameters as per product label claim:

1. Directions
2. Storage
3. Expiry

Warranty

This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

Reference

1. Koneman E. W., Allen S. D., Janda W.M., Schreckenberger P.C., Winn W.C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company
2. Vera H. D., 1950, Am. J. Public Health, 40, 1267
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification –Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
4. Data on file: Microxpress®, A Division of Tulip Diagnostics (P) Ltd.

Product Presentation:

Cat No.	Product description	Pack Size
201160160100	Dehydrated Culture Media	100 g

	Temperature Limit		Manufacturer		Batch Code		Date of Manufacture		This way up		Received on
	Catalogue Number		Consult Instructions for use		Use-by Date		Hygroscopic keep container tightly closed		Opened on		

Revision: 0825/VER-03

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.