

Taurocholate Broth**Intended Use**

Taurocholate Broth is used for selective isolation of coliforms from water, milk and other food products.

Summary

Taurocholate Broth selectively supports growth of Gram-negative coliforms from water, milk and other similar food products.

Principle

The selective action of this medium is attributed to crystal violet and sodium taurocholate which inhibits most species of Gram-positive bacteria. Lactose is a fermentable carbohydrate. Neutral red is the indicator dye in the media. Lactose fermenters acquire pinkish red colour due to production of acid. Sodium chloride maintains the osmotic balance at the medium.

Formula*

Ingredients	g/L
Casein Enzymic Hydrolysate	8.0
Peptic Digest of Animal Tissue	12.0
Lactose	10.0
Sodium Taurocholate	5.0
Sodium Chloride	5.0
Neutral Red	0.03
Crystal Violet	0.0001
Final pH (at 25°C)	7.4 ± 0.2

*Adjusted to suit performance parameters.

Type of Specimen

Food and dairy samples; Water samples

Storage and Stability

Store dehydrated medium below 30°C in tightly closed container and the prepared medium at 2°C-8°C. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Specimen Collection and Handling

Ensure that all samples are properly labelled.

Follow appropriate techniques for handling samples as per established guidelines.

Some samples may require special handling, such as immediate refrigeration or protection from light, follow the standard procedure.

The samples must be stored and tested within the permissible time duration.

After use, contaminated materials must be sterilized by autoclaving before discarding.

Directions

1. Suspend 40.03 g of the powder in 1000 mL purified / distilled water.
2. Heat if necessary, to dissolve the powder completely.
3. Dispense in tubes containing inverted and Durham's tubes and sterilize by autoclaving at 121°C (15 psi) for 15 minutes as per validated cycle.

Quality Control

Dehydrated Appearance: Light yellow to pink coloured, homogeneous, free flowing powder.

Prepared Appearance: Red coloured, clear solution without any precipitate.

Cultural Response: Cultural characteristics observed after an incubation of 18-48 hours at 30°C-35°C.

Organism (ATCC)	Growth	Acid	Gas
<i>Klebsiella aerogenes</i> (13048)	Good	+	+
<i>Escherichia coli</i> (25922)	Good	+	+
<i>Escherichia coli</i> (8739)	Good	+	+
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> (25923)	Inhibited	—	—
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> (6538)	Inhibited	—	—

Performance and Evaluation

Performance of the product is dependent on following parameters as per product label claim:

1. Directions
2. Storage
3. Expiry

Warranty

This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

Reference

1. Data on file: Microxpress®, A division of Tulip Diagnostics (P) Ltd.

Product Presentation:

Cat No.	Product description	Pack Size
201200040100	Dehydrated Culture Media	100 g
201200040500	Dehydrated Culture Media	500 g

 Temperature Limit	 Manufacturer	 LOT Batch Code	 Date of Manufacture	 This way up	 RO Received on
REF Catalogue Number	 Consult Instructions for use	 Use-by Date	 Hygroscopic keep container tightly closed	 OO Opened on	

Revision: 0825/VER-03

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.