Tryptone Soya Yeast Extract Agar ISO

Intended Use

Tryptone Soya Yeast Extract Agar ISO is used for confirmation of *Listeria* in Henry's light in compliance with ISO specifications ISO 10560:1993.

Summary

Tryptone Soya Yeast Extract Agar is formulated as per APHA for the isolation and cultivation of *L. monocytogenes* from foods. ISO Committee has recommended this medium for confirmation of *Listeria* species and can also be used for the cultivation and maintenance of a wide variety of heterotrophic microorganisms.

Principle

Casein enzymic hydrolysate and papaic digest of soyabean meal provide amino acids and other complex nitrogenous substances. Dextrose is the energy source. Dipotassium hydrogen phosphate buffers the medium. Yeast extract is the rich source of vitamin B complex. Agar is a gelling agent.

Formula*

Ingredients	g/L
Casein Enzymic Hydrolysate	17.0
Papaic Digest of Soyabean Meal	3.0
Sodium Chloride	5.0
Dipotassium Hydrogen Phosphate	2.5
Dextrose	2.5
Yeast Extract	6.0
Agar	15.0
Final pH (at 25°C)	7.3 ± 0.2
*Adjusted to suit performance parameters.	

Storage and Stability

Store dehydrated medium below 30°C in tightly closed container and the prepared medium at 2°C-8°C. Avoidfreezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Type of Specimen

Pharmaceutical samples and Food Samples.

Specimen Collection and Handling

Ensure that all samples are properly labelled.

Follow appropriate techniques for handling samples as per established guidelines. Some samples may require special handling, such as immediate refrigeration or protection from light, follow the standard procedure. The samples must be stored and tested within the permissible time duration. After use, contaminated materials must be sterilized by autoclaving before discarding.

Directions

- 1. Suspend 51.0 g of the powder in 1000mL purified / distilled water.
- 2. Heat if necessary, to dissolve the powder completely.
- 3. Sterilize by autoclaving at 121°C (15 psi) for 15 minutes.
- 4. Mix well and pour into sterile petridishes.

Quality Control

Dehydrated Appearance: Cream to yellow coloured, homogenous, free flowing powder.

Prepared Appearance: Light amber coloured, clear to slightly opalescent gel forms in petridishes.

Growth Promotion Test: Growth promotion is carried out in accordance with the harmonized method of USP/EP/JP/IP and growth is observed after an incubation at 30-35°C for 18 to 24 hours.

Growth Promoting Properties: The test results observed are within the specified temperature and shortest period of time specified in the test, inoculating \leq 100 cfu of appropriate microorganism at 30-35°C for 18 hours.

Organism (ATCC)	Growth
Listeria monocytogenes strain Li 23 (19114)	Good
Listeria monocytogenes serotype 4b (19115)	Good

Note: For Good growth - Growth obtained on test media should not differ by a factor greater than 2 from calculated value for a standardized inoculum.

Performance and Evaluation

Performance of the product is dependent on following parameters as per product label claim:

- 1. Directions
- 2. Storage
- 3. Expiry

Warranty

This product is designed to perform as described on the label and package insert. The manufacturer disclaims any implied warranty of use and sale for any other purpose.

References

1. Vanderzant C. and Splittstoesser D. F., (Eds.), 1992, Compendium of Methods for the MicrobiologicalExamination of Foods, 3rd Ed., APHA, Washington, D.C.

Pack Size

- 2. International Organization for Standardization (ISO), 1993, Draft, ISO/DIS 10560.
- 3. Atlas R. M. 2004, 3rd Ed., Handbook of Microbiological Media, Parks, L.C. (Ed.), CRC Press, Boca Raton.
- 4. FDA, Bacteriological Analytical Manual, 2005, 18th Ed., AOAC, Washington, DC.
- 5. Data on file: Microxpress[®], A Division of Tulip Diagnostics (P) Ltd.

Product Presentation:

Cat No.	Product description
201200320500	Dehydrated Culture Media

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.