

Yeast Peptone

Intended Use

A nutritious extract used in the preparation of culture media for the cultivation of a wide variety of microorganisms.

Summary and Principle

Yeast Peptone is obtained by enzymatic hydrolysis of yeast protein. It can be used as replacement for animal and vegetable origin peptones. Yeast peptone is used in fermentation industries such as Probiotics, and also in Biopharmaceuticals.

Storage and Stability

Store dehydrated medium below 30°C in tightly closed container. Avoid freezing and overheating. Use before expiry date on the label. Once opened keep powdered medium closed to avoid hydration.

Note: TSE/BSE certificate is available on request.

Directions

Refer to the final concentration in the formula of the medium being prepared.

Quality Control

Test	Specification
Appearance	Light yellow coloured powder.
Solubility	Completely soluble in water.
Colour and Clarity of 1% w/v aqueous solution after autoclaving at 15 psi / 15 min	Light yellow coloured, clear solution.
pH after autoclaving	6.0 ± 0.5
Ash Content	Not More Than 15%
Loss on Drying (Moisture Content)	Not More Than 5%
α-amino Nitrogen Content	Not Less Than 3%
Total Nitrogen Content	Not Less Than 15%
Total microbial count	Less than 5000 cfu/g
<i>E. coli</i>	Absent
<i>Salmonella</i>	Absent
<i>Pseudomonas aeruginosa</i>	Absent
<i>Staphylococcus aureus</i>	Absent

Cultural Response

Cultural characteristics observed after an incubation of 18-24 hours at 30°C-35°C for bacteria and 2-5 days for fungi at 20°C-25°C

Organism (ATCC)	Growth
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> (6538)	Good
<i>Escherichia coli</i> (8739)	Good
<i>Pseudomonas aeruginosa</i> (9027)	Good
<i>Streptococcus pyogenes</i> Strain Bruno (19615)	Good
<i>Candida albicans</i> 3147 (10231)	Good
<i>Aspergillus brasiliensis</i> WLRI 034(120) (16404)	Good











Note: Growth for *Aspergillus brasiliensis* was observed after 72 hours at 20°C-25°C for quantitative test and the same is carried out for qualitative test and confirmed characteristic growth (White mycelial growth with black spores) after 4-5 days.

Reference

1. Data on file: Microxpress®, A Division of Tulip Diagnostics (P) Ltd.

Product Presentation:

Cat No.	Product description	Pack Size
202250140500	Yeast Peptone	500 g
202250142500	Yeast Peptone	2.5 k
202250149925	Yeast Peptone	25 k (Bag)
202250149825	Yeast Peptone	25 k (Drum)

									
Temperature Limit	Manufacturer	Lot Number	Hygroscopic keep container tightly closed	Date of Manufacture	Catalogue Number	Consult Instructions for use	Use-by Date	Received on	Opened on

Revision: 1025/VER-03

Disclaimer

Information provided is based on our inhouse technical data on file, it is recommended that user should validate at his end for suitable use of the product.
